

Styrene Expert Panel Report

Part B — Recommendation for listing status for “styrene” in the Report on Carcinogens and scientific justification for the recommendation

The Report on Carcinogens (RoC) expert panel for styrene met at the Radisson Hotel, Durham, North Carolina on July 21-22, 2008, to peer review the draft background document on styrene and make a recommendation for the listing status in the 12th Edition of the RoC. Members of the expert panel are as follows:

Phillips, David, Ph.D., D.Sc., (Chair) Institute of Cancer Research	Snedeker, Suzanne, Ph.D. Program on Breast Cancer and Environmental Risk Factors College of Veterinary Medicine Cornell University
Eustis, Scott, D.V.M., Ph.D., DACVP Retired, formerly Senior Director, Regulatory Pathology Department of Safety Sciences, Pfizer	Stone, Michael, Ph.D. Department of Chemistry Vanderbilt University School of Medicine.
Infante, Peter, Dr.P.H., M.P.H., D.D.S Peter Infante Consulting, LLC	Ward, Elizabeth M., Ph.D. Surveillance Research Epidemiology and Surveillance Research American Cancer Society
Matanoski, Genevieve, M.D., Dr.P.H. Department of Epidemiology Johns Hopkins Bloomberg School of Public Health	Yost, Garold S., Ph.D. Department of Pharmacology and Toxicology University of Utah
Que Hee, Shane S., Ph.D UCLA School of Public Health Environmental Health Sciences Center for Occupational and Environmental Health	Zeise, Lauren, Ph.D. California EPA OEHHA Reproductive & Cancer Hazard Assessment
Smith, Thomas J., Ph.D., CIH Department of Environment Health Harvard School of Public Health	

The recommendation for listing status and scientific justification for the recommendation follow this page.

Overall Evaluation

Following a discussion of the draft document, together with additional recommended inclusions and some re-analysis of published data, the expert panel reviewed the RoC listing criteria and made its recommendation for the listing status of styrene in the RoC. The expert panel recommended by a vote of 8 yes/2 no that styrene should be listed in the RoC as *reasonably anticipated to be a human carcinogen* based on limited evidence of carcinogenicity in humans and sufficient evidence in animals. The two members of the panel who voted against the motion did so because, in their opinion, styrene should be listed as *known to be a human carcinogen*.

The major considerations discussed by the panel that led to its majority recommendation included: (1) evidence of past and present human exposure to styrene in the United States; (2) evidence of cancer in styrene-exposed workers; (3) induction of lung tumors in mice by styrene by two routes of exposure; (4) the established carcinogenicity in animals and genotoxicity of a styrene metabolite, the 7,8-oxide; (5) evidence for styrene-related DNA adducts and cytogenetic effects in styrene-exposed workers.

This evidence is described in further detail below.

Human Exposure

There is more than sufficient data to show that humans are exposed to styrene by inhalation, skin exposure, and oral absorption in such situations as the workplace (inhalation and skin exposure predominantly), food and drinking water (oral ingestion), and in the ambient environment (levels found in ambient air, surface and groundwater, and soils) in the United States.

Human Cancer Studies

The expert panel concluded that there was limited evidence for the carcinogenicity of styrene in humans, although some members were of the opinion that the epidemiological data provided sufficient evidence. The strongest evidence for cancer in humans is the association between styrene exposure and non-Hodgkin lymphoma (NHL). This evidence comes from the Delzell *et al.* (2006) analysis in the styrene-butadiene industry and the Kogevinas (1994a) study in the reinforced plastics industry. In the Delzell study there was an exposure-response relationship for NHL and NHL plus chronic lymphocytic leukemia (CLL) that was not attenuated by control for butadiene and only mildly attenuated by control for dimethyldithiocarbamate (DMDTC) (which may not have been appropriate to control for). It is very unlikely that such a strong exposure-response trend could be due to chance, bias, or confounding. These findings are supported by increases in RR for all lymphomas with time since first exposure and estimated average exposure in the multiplant-cohort studied by Kogevinas. Many of the smaller studies found excesses in lymphohematopoietic cancers, although these were not statistically significant. They provide further evidence of consistency in the risk. In addition, there is some evidence for increased risk of pancreatic and esophageal cancer. It should be noted that increased risk of solid tumors was not evaluated in many studies. Studies in workers have provided evidence for DNA adducts and chromosomal aberrations in lymphocytes of styrene-exposed workers which supports a genotoxic mechanism for styrene.

Studies on Experimental Animals

Evidence on the carcinogenicity of styrene comes from a number of studies in rats and mice. There are robust studies in male and female mice and rats by oral gavage (NCI 1979a) and inhalation (Cruzan *et al.* 2001, Cruzan *et al.* 1998) routes, along with several other studies more limited in their ability to detect carcinogenic effects because of study design (low doses, short treatment, short study duration, small group size), high early mortality, or limited reporting (tumor diagnosis) (Beliles *et al.* 1985, Brunnemann *et al.* 1992, Conti *et al.* 1988, Jersey *et al.* 1978, Ponomarev and Tomatis 1978).

Sufficient evidence of carcinogenic activity comes from multiple studies in mice exposed to styrene by multiple routes. Styrene induced benign and malignant lung tumors in male and female mice by inhalation (Cruzan *et al.* 2001) and in male mice by oral intubation (NCI 1979a). This is supported by findings of lung tumors in both sexes of mice in studies of more limited design (Ponomarev and Tomatis 1978). There is also the finding in rats of malignant mammary tumors by inhalation (Conti *et al.* 1988) and a small increase in mammary fibroadenoma in a relatively low dose drinking water study (Beliles *et al.* 1985), but mammary tumors were not increased in an adequate inhalation study in the same rat strain exposed for two years (Cruzan *et al.* 1998), limiting the weight given to the mammary tumor findings. In earlier reviews (IARC 1994b, NTP 2002), sufficient evidence in animals was found for the carcinogenicity of styrene-7,8-oxide.

Mechanistic Concerns

There are at least two reasonable, literature-supported, and scientifically valid mechanisms for styrene-induced cancer. These mechanisms are not mutually exclusive. These mechanisms supported the panel's decision to recommend listing styrene in the Report on Carcinogens.

- Styrene is genotoxic to human lymphocytes through sister chromatid exchange processes and chromosomal aberrations (e.g., micronuclei and aneuploidy). Styrene is mutagenic through formation of styrene-7,8-oxide. Cyp2f2 in mouse lung cells efficiently oxidizes styrene to styrene-7,8-oxide. In addition, CYP2F1, CYP2A13, and CYP2S1 are expressed in many extrahepatic organs, and their known catalytic activity with styrene supports a plausible mechanism by which styrene is bioactivated to styrene oxide in many human tissues. Styrene-7,8-oxide forms multiple DNA adducts, primarily O⁶ and N7 of guanine, in cultured human lymphocytes and embryonal lung fibroblasts. In addition, lymphocytes from styrene-exposed workers possess the same DNA adducts.
- Styrene bioactivation to the 7,8-epoxide or 3,4-epoxide is cytotoxic to mouse lung cells. Specifically, the 3,4-epoxide could form 4-vinylphenol which could be oxidized to dihydrodiols and quinones by P450 enzymes and produce cytotoxicity through reactive oxygen mechanisms. Subsequent cellular proliferation and clonal expansion would then constitute an epigenetic mechanism. Recent evidence suggests that mice may be especially sensitive to lung carcinogenesis through this mechanism, and it may be operative in humans as well.

Organ Selectivity

- Styrene is potentially carcinogenic to lung tissues because sufficient evidence exists in animals for bioactivation to the mutagenic epoxide, and tumors are produced within the same sites where P450s are expressed, and DNA adducts are also identified.
- Support for a mechanism of mammary tissue as a site includes limited human evidence for elevated prolactin levels from styrene-exposed workers.
- Styrene produces lymphohematopoietic cancer in humans, and this effect has the strongest epidemiological support. However, bioactivation of styrene to styrene-7,8-oxide by cytochrome P450 enzymes in blood cells has not been reported. Thus, carcinogenic mechanisms for lymphomas would likely depend on direct interactions of preformed styrene-7,8-oxide in blood cells.

Report Approved

Redacted

David Phillips, Ph.D. (Chair)

Date

Aug 26 2008

References

1. Beliles RP, Butala JH, Stack CR, Makris S. 1985. Chronic toxicity and three-generation reproduction study of styrene monomer in the drinking water of rats. *Fundam Appl Toxicol* 5(5): 855-868. (Supported by the Chemical Manufacturers Association Styrene Program Panel. Authors affiliated with Litton Bionetics, Inc., USA; Gulf Oil Corporation, USA; Chemical Manufacturers Association, USA; George Washington University, USA.)
2. Brunnemann KD, Rivenson A, Cheng SC, Saa V, Hoffmann D. 1992. A study of tobacco carcinogenesis. XLVII. Bioassays of vinylpyridines for genotoxicity and for tumorigenicity in A/J mice. *Cancer Lett* 65(2): 107-113. (Supported by the U.S. National Cancer Institute and the American Cancer Society. Authors affiliated with the American Health Foundation, NY.)
3. Conti B, Maltoni C, Perino G, Ciliberti A. 1988. Long-Term Carcinogenicity Bioassays on Styrene Administered by Inhalation, Ingestion and Injection and Styrene Oxide Administered by Ingestion in Sprague-Dawley Rats, and *para*-Methylstyrene Administered by Ingestion in Sprague-Dawley Rats and Swiss Mice. In *Living in a Chemical World*, Annals of the New York Academy of Sciences, vol. 534. Maltoni C, Selikoff IJ, eds. New York, NY: New York Academy of Sciences. pp. 203-234. (Support not reported. Authors affiliated with Institute of Oncology "F. Addarii", Italy.)
4. Cruzan G, Cushman JR, Andrews LS, Granville GC, Johnson KA, Hardy CJ, Coombs DW, Mullins PA, Brown WR. 1998. Chronic toxicity/oncogenicity study of styrene in CD rats by inhalation exposure for 104 weeks. *Toxicol Sci* 46(2): 266-281. (Supported by Styrene Information and Research Center. Authors affiliated with ToxWorks, NJ; Chevron Research and Technology, CA; ARCO Chemical Company, PA; Shell Canada Ltd., Canada; Dow Chemical Company, MI; Huntingdon Research Center, UK; Research Pathology Services, PA.)
5. Cruzan G, Cushman JR, Andrews LS, Granville GC, Johnson KA, Bevan C, Hardy CJ, Coombs DW, Mullins PA, Brown WR. 2001. Chronic toxicity/oncogenicity study of styrene in CD-1 mice by inhalation exposure for 104 weeks. *J Appl Toxicol* 21(3): 185-198. (Supported by the Styrene Information and Research Center. Authors affiliated with ToxWorks, NJ; Chevron Research and Technology, CA; Rohm & Hass Company, PA; Shell Canada Ltd., Canada; Dow Chemical Company, MI; Huntingdon Research Center, UK; BP Amoco Chemical Co., IL; Research Pathology Services, PA.)
6. Delzell E, Sathiakumar N, Graff J, Macaluso M, Maldonado G, Matthews R. 2006. An updated study of mortality among North American synthetic rubber industry workers. *Res Rep Health Eff Inst*(132): 1-63; discussion 65-74.
7. IARC. 1994b. *Styrene Oxide*. Volume 36. IARC Monographs on the Evaluation of Carcinogenic Risk of Chemicals to Humans. Lyon, France: International Agency for Research on Cancer.
8. Jersey GC, Balmer MF, Quast JF, Park CN, Schuetz DJ, Beyer JE, Olson KJ, McCollister SB, Rampy LW. 1978. *Two-Year Chronic Inhalation Toxicity and Carcinogenicity Study*

on Monomeric Styrene in Rats. Final Report. Michigan: Toxicology Research Laboratory, Health and Environmental Research, Dow Chemical U.S.A.

9. Kogevinas M, Ferro G, Andersen A, Bellander T, Biocca M, Coggon D, Gennaro V, Hutchings S, Kolstad H, Lundberg I, Lynge E, Partanen T, Saracci R. 1994a. Cancer mortality in a historical cohort study of workers exposed to styrene. *Scand J Work Environ Health* 20(4): 251-261. (Supported by the Commission of the European Communities. Authors affiliated with IARC; Cancer Registry of Norway; Karolinska Hospital, Sweden; Presidio Multizonale di Prevenzione, Italy; University of Southampton, UK; Istituto Nazionale per la ricerca sul cancro, Italy; Health and Safety Executive, UK; University of Aarhus, Denmark; Danish Cancer Registry; Institute of Occupational Health, Finland.)
10. NCI. 1979a. *Bioassay of Styrene for Possible Carcinogenicity.* Technical Report Series No. 185. Bethesda, MD: National Cancer Institute.
11. NTP. 2002. *Report on Carcinogens Background Document for Styrene-7,8-oxide*, Research Triangle Park, NC: National Toxicology Program. 144 pp. <http://ntp-server.niehs.nih.gov/ntp/newhomeroc/roc10/SO.pdf>.
12. Ponomarev V, Tomatis L. 1978. Effects of long-term oral administration of styrene to mice and rats. *Scand J Work Environ Health* 4(Suppl 2): 127-135. (Support not reported. Authors affiliated with IARC, France.)